



KILCOX EXTRA



A bactericidal, yeasticidal and virucidal product for the disinfection of livestock premises, animal breeding equipment and materials and the disinfection of boots in farms.









KILCOX EXTRA



ADVANTAGES

- Broad spectrum disinfectant based on QAC, Glutaral and Chlorocresol
- For surfaces, equipment, materials in animal houses or farms and boot disinfection
- Bactericidal, yeasticidal, virucidal efficacy
- Tested Efficacy against Avian Influenza (AI) Virus













Surface & Equipment Disinfection in livestock housing

Spray application at low pressure:

Pre-clean surfaces and equipment to be disinfected. Apply the diluted product by spraying at low pressure at an average rate of 300 ml/m² up to the limit of runoff at 10°C minimum:

• Bactericidal: 0.5% / ≥ 30 minutes contact time

Virucidal: 2% / ≥ 30 minutes contact time

• Yeasticidal: 3.5% / ≥ 60 minutes contact time

Equipment Disinfection / Boot Disinfection

Application by dipping / boot bath application

Pre-clean equipment, materials and boots to be disinfected. Prepare a dilution of KILCOX EXTRA at the recommended rate and apply it by dipping in equipment, materials or boots at a minimum of 10°C:

• Bactericidal: 0.5% / ≥ 30 minutes contact time

• Virucidal: 2% / ≥ 30 minutes contact time

Yeasticidal: 3.5% / ≥ 60 minutes contact time

BACTERICIDAL EFFICACY

EN 14349 / 0.5% / 30 MIN. / 10°C / high soiling (Enterococcus hirae, Proteus vulgaris, Pseudomonas aeruginosa, Staphylococcus aureus)

EN 1656 / 0.5% / 5 MIN. / 10°C / low soiling (Enterococcus hirae, Proteus vulgaris, Pseudomonas aeruginosa, Staphylococcus aureus)

YEASTICIDAL EFFICACY

EN 16438 / 3.5% / 60 min. / 10°C / high soiling (Candida albicans)

EN 1657 / 1% / 30 min. / 10°C / high soiling (Candida albicans)

Active substance: QAC, Glutaral, Chlorocresol

Storage: Keep in a clean, cool and well-ventilated place away from sources of heat and

intense light.

Shelf Life: 24 months from date of manufacture.

VIRUCIDAL EFFICACY

EN 14675 / 2% / 30 MIN. / 10°C / low soiling (Bovine Enterovirus type 1 (ECBO)) (Avian Influenza Virus (H5N2, H1N1)

DEFRA:

At a dilution rate of 1 ml disinfectant plus 110 ml water
Influenza of Avian Origin in Mammals
At a dilution rate of 1 ml disinfectant plus 300 ml water
Swine Vesicular Disease

EFFICACY - DEFRA

At a dilution rate of 1 ml disinfectant plus 50 ml water General orders

