

VIROPHOR 2.8%



Pig &
Poultry
Farming

**An iodine-based
disinfectant suitable for
disinfection of foot wear
and surfaces in animal
and poultry housing.**



VIROPHOR 2.8%



ADVANTAGES

- Broad spectrum disinfectant based on Iodine
- Bactericidal, Virucidal and Yeasticidal efficacy
- For surfaces in animal houses or farms
- For disinfection of boot / footwear immersion



Spray



Foam

Surface Disinfection in livestock housing

Spray and Foam application at low pressure – only by trained personnel:

Pre-clean surfaces and equipment to be disinfected. Apply the diluted product by spraying at low pressure or foaming at an average rate of 300 ml/m² up to the limit of runoff at 10°C minimum:

- Bactericidal & yeasticidal: 2% / ≥ 1 minute contact time
- Virucidal: 2% / ≥ 30 minutes contact time



Dip

Footwear Disinfection / Boot Disinfection

Application by dipping / boot bath application – only by trained personnel:

Pre-clean boots to be disinfected. Prepare on a daily base a dilution of VIROPHOR 2.8% at the recommended rate and apply it by dipping boots at a min. of 10°C:

- Bactericidal & Yeasticidal: 2% / ≥ 1 minute contact time

Keep foot dip covered when not in use. Specific instruction that this product is for use as a boot dip under supervised conditions only. It is prohibited from use as a general boot scrub/ disinfectant.

BACTERICIDAL EFFICACY

EN 14349 & EN 1656 / 2% / 1 MIN. / 10°C / low soiling
(*Enterococcus hirae*, *Proteus vulgaris*, *Pseudomonas aeruginosa*,
Staphylococcus aureus)

YEASTICIDAL EFFICACY

EN 16438 & EN 1657 / 2% / 1 min. / 10°C / low soiling
(*Candida albicans*)

VIRUCIDAL EFFICACY

EN 14675 / 2% / 30 MIN. / 10°C / low soiling
(*Bovine Enterovirus type 1 (ECBO)*)

EFFICACY – DEFRA

At a dilution rate of 1 ml disinfectant plus 550 ml water
Foot and Mouth Disease: 1:550

At a dilution rate of 1 ml disinfectant plus 150 ml water
Swine vesicular Disease: 1:150

At a dilution rate of 1 ml disinfectant plus 99 ml water
Diseases of Poultry Order (Newcastle): 1:99
Avian Influenza Order: 1:99

Influenza of Avian Origin in Mammals: 1:99

At a dilution rate of 1 ml disinfectant plus 50 ml water
General orders

At a dilution rate of 1 ml disinfectant plus 20 ml water
Tuberculosis: 1:20

Active substance: Iodine

Caution: To be used by trained professionals - trained in the use of this product and also with respect to the specified PPE/RPE. Due to the corrosive nature of the concentrate and diluted product it is not suitable for general use by untrained farm operatives.

Risk mitigation: Feed, water and troughs are to be covered during biocidal treatment. Wash hands thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection.

Storage: Keep in a clean, cool and well-ventilated place away from sources of heat and intense light. Store in tightly closed original container. Protect from frost. Store below 25°C and out of direct sunlight.

Shelf Life: 24 months from date of manufacture.